

# Evaluation of Bacteriological Parameters of Water Quality in The Bouregreg Estuary Along the Moroccan Atlantic Coast

M. El Amraoui<sup>1,2</sup>, M. Tarbaoui<sup>3,\*</sup>, J. Benba<sup>4</sup>, B. El Amraoui<sup>1,5</sup>, T. Bamhaoud<sup>1</sup>,  
N. Lazrak<sup>2</sup>, M. Monkade<sup>6</sup>

**Abstract-** On the Moroccan Atlantic coast, the Bouregreg estuary was submitted to many anthropic effects. After the dam Sidi Mohammed ben Abdellah was built in 1974, the flow became almost null. The waters of the estuary are meant for recreational activities, fishing, but at the same time they receive water collecting domestic and industrial wastewater. The waste in old landfills was not treated and leaked into the river water. The matter gets worse with the action of tides which make it difficult to escape the pollution load (physicochemical, bacteriological, heavy metals, etc.). The main objective of this study is to monitor the bacteriological quality of the surface waters from Bouregreg estuary (Moroccan Atlantic coast). The bacteriological parameters were determined upstream (S1, S2 and S3) and downstream (S4, S5 and S6) of the Bouregreg estuary. The parameters studied are fecal coliform (FC), fecal streptococci (FS), Salmonella, Staphylococcus aureus and Pseudomonas aeruginosa. The evaluation of the bacterial contamination of the studied water samples shows that the rate of contamination by fecal coliforms is very high in station S3 compared to other stations. This is explained by the emplacement of this station which is located next to a main rejection of Oued Akrach. The bacterial loads are maximal in this station:  $7 \cdot 10^3$  CF/100ml for fecal coliforms and  $2 \cdot 10^3$  SF/100ml for fecal streptococci. The ratio values (CF/SF) for the different stations show that the fecal pollution observed at the studied estuary is of mixed origin with human dominance. The search for pathogenic bacteria namely Salmonella, Staphylococcus aureus and Pseudomonas aeruginosa, was negative for all the studied samples.

**Index Terms**— Atlantic coast; Bacteriological parameters; Bouregreg estuary; Morocco; Pollution; Water quality.

Mohammed El Amraoui,<sup>1</sup>Laboratoire Contrôle Qualité en Bioindustrie et Molécules Bioactives, Faculté des Sciences, Université Chouaib Doukkali, BP 20, 24000 El Jadida, Maroc, <sup>2</sup>Laboratoire du Maghreb, BP 2863 Rabat, Maroc

Meriem Tarbaoui,<sup>3</sup>Laboratoire Biomolécules et Synthèse Organique, Faculté des Sciences Ben M'sik, Université Hassan II-Casablanca, B.P 7955 Casablanca, Maroc

Jamila Benba,<sup>4</sup>laboratoires de Biotechnologie Marine et de l'Environnement, Faculté des Sciences, Université Chouaib Doukkali, BP 20, 24000 El Jadida, Maroc

Belkassem El Amraoui,<sup>1</sup>Laboratoire Contrôle Qualité en Bioindustrie et Molécules Bioactives, Faculté des Sciences, Université Chouaib Doukkali, BP 20, 24000 El Jadida, Maroc, <sup>5</sup>Physicochimie des Milieux Naturels et Molécules Bioactives (PMNMBA), Faculté Polydisciplinaire de Taroudant, Université Ibn Zohr, BP 271, 83000 Taroudant, Maroc,

Toufiq Bamhaoud,<sup>1</sup>Laboratoire Contrôle Qualité en Bioindustrie et Molécules Bioactives, Faculté des Sciences, Université Chouaib Doukkali, BP 20, 24000 El Jadida, Maroc, <sup>2</sup>Laboratoire du Maghreb, BP 2863 Rabat, Maroc

Nicole Lazrak,<sup>2</sup>Laboratoire du Maghreb, BP 2863 Rabat, Maroc

Mohammed Monkade,<sup>6</sup>Département de Physique, Faculté des Sciences, Université Chouaib Doukkali, BP 20, 24000 El Jadida, Maroc .

## I. INTRODUCTION

An estuary is the place of transition between salt water and fresh water. The historical importance of these geographical areas is due to the existing shops and to the shipping of products. On the Moroccan Atlantic coast, the Bouregreg estuary was submitted to many anthropic effects. After the dam Sidi Mohammed Ben Abdellah was built in 1974, the flow became almost null. The estuary is bordered by two cities, Rabat and Sale in which the process of urbanization and industrialization has continually increased, affecting the water quality. The waters of the estuary are meant for recreational activities, fishing, but at the same time they receive water collecting domestic and industrial wastewater. The waste in old landfills was not treated and leaked into the river water. The matter gets worse with the action of tides which make it difficult to escape the pollution load (physicochemical, bacteriological, heavy metals, etc.). Knowing that the fishing sites and shellfish gathering sites are close to discharge points, we can imagine how serious the consequences could be. Biological monitoring is essential in assessing the quality of the estuary water. In Moroccan estuaries, several authors have shown such contamination [1] -[2]- [3]- [4] due to domestic and industrial activities along the Moroccan coast. The objective of this study is to evaluate the bacteriological quality of the water of the Bouregreg estuary and to predict their impact on the global state of the environment and the state of the marine ecosystems in particular.

## II. MATERIALS AND METHODS

### A. Presentation of the study area (Bouregreg estuary)

The Bouregreg estuary is located on the Atlantic between the two cities Rabat and Sale  $34^\circ$  N and  $6^\circ 50'$  west. It has a length of 23 km, limited by the dam of Sidi Mohammed Ben Abdellah and an average width of 150 m. It is generally facing south-east and north-west, except in the area ranging from kilometer point (kp) 13.5 to the confluence of Oued Akrech where it is oriented sud-ouest/north – East. Upstream and up the river Akrach (18 km), the estuary has an appearance of deep valley surrounded by highlands; Downstream of kilometer point (kp) 14.5 (Both islands), the collection disappears and the estuary through an alluvial plain in which he describes many meanders. This terrace, called Oulja disappears, about 4 km from the ocean where marshes are observed. Near its mouth are two sandy beaches of Rabat and



➤ The pre-enrichment

On the first day, 5 liters of water to be analyzed is filtered and the filtration membranes are placed in a flask of buffered peptone water and incubated at 37 °C for 24 hours to revive the stressed bacteria.

➤ The enrichment

On the second day, 1 ml of the buffered peptone water is placed in a Selenite and GN broth and then incubated at 44 °C for 24 hours.

➤ Isolation

On the third day, the isolation is carried out from selenite and GN on Hektoen medium.

Salmonella give black colonies with a greenish halo on Hektoen.

➤ Enumeration of germs

For salmonella we can only use the method of presence or absence, we cannot make an enumeration since we have made enrichment.

*Staphylococcus aureus*

Staphylococcus aureus is a gram-positive coccil bacterium that is a member of the Firmicutes, and is frequently found in the nose, respiratory tract, and on the skin. It is often positive for catalase and nitrate reduction and is a facultative aerobe that can grow without the need for oxygen. [14]

Although S. aureus is not always pathogenic, it is a common cause of skin infections such as abscesses, respiratory infections such as sinusitis, and food poisoning. Pathogenic strains often promote infections by producing potent protein toxins, and expressing cell-surface proteins that bind and inactivate antibodies.

*Pseudomonas aeruginosa*

Pseudomonas aeruginosa is a bacterium found frequently in some water systems and in the environment. Its origin may be human and possibly fecal. Its presence in groundwater can result in contamination by surface water. It is found frequently in wastewater with concentrations up to 106/100 ml. It causes infections of the ears, eyes, burns and urinary tract as well as enteritis.

*Identification by ATB EXPRESSION*

To confirm the identification of the germs found in water samples, the identification by ATB EXPRESSION was used. It is a semi-automated biochemical identification of which the identification galleries containing the dehydrated substrates are inoculated with a standardized bacterial suspension and then incubated for 4 or 24 hours at a given temperature. After incubation, the galleries are read by the ATB identification system and then the data are interpreted.

III. RESULTS AND DISCUSSIONS

A. Evaluation of the degree of fecal contamination in the different sampling points

The results of the bacteriological analysis of water from six studied stations reveal the presence of the germs of fecal contamination (Table I). The fecal coliform load average (FC) varies between 2.6 10<sup>3</sup> and 7 10<sup>3</sup> (CFU / 100 mL). In terms of bacterial load, the station S3 is more concentrated with fecal staphylococci than other stations. The fecal

staphylococci load average (FS) varies between 0.9 10<sup>3</sup> and 2 10<sup>3</sup> (CFU / 100 mL). In terms of bacterial load, the stations S3 and S4 are slightly more concentrated with fecal staphylococci than other stations. The result of the enumeration of the germs of fecal contamination is in good agreement with the bibliographical data about the state of bacterial contamination of urban effluents [15]- [17], but it far exceeds the standard set by the World Health Organization 1000 CFU / 100 mL [18]. The ratio values (FC/FS) for the different studied stations show that the fecal pollution observed at the studied estuary is of mixed origin with human dominance (Table II).

Table I. The average values of bacteriological parameters of water samples in the six studied stations

Parameters Stations	Fecal coliform (CFU /100 mL)	Fecal staphylococci (CFU /100 mL)
Station S1	3 10 <sup>3</sup>	1 10 <sup>3</sup>
Station S2	2.6 10 <sup>3</sup>	0.9 10 <sup>3</sup>
Station S3	7 10 <sup>3</sup>	2 10 <sup>3</sup>
Station S4	3.7 10 <sup>3</sup>	2 10 <sup>3</sup>
Station S5	4 10 <sup>3</sup>	1.1 10 <sup>3</sup>
Station S6	3.2 10 <sup>3</sup>	1.2 10 <sup>3</sup>

Table II. Origin of contamination in the six studied stations

Sampling stations	Report (FC/FS)	Origin of contamination
Station S1	3	Mixed with human dominance
Station S2	2.88	Mixed with human dominance
Station S3	3.5	Mixed with human dominance
Station S4	1.85	Uncertain
Station S5	3.63	Mixed with human dominance
Station S6	2.6	Mixed with human dominance

B. Evaluation of the degree of contamination of the different sampling points by pathogenic germs

After having used all the biochemical tests necessary for the identification of Salmonella, none was found to be salmonella for all the water samples studied. These results are in agreement with some anterior studies [19]. The search of staphylococcus aureus and pseudomonas aeruginosa was also negative for all the water samples studied. Several causes can be at the origin of the absence of the pathogenic germs of which the most important are the high elevated values of the pH and the concentrations of oxygen dissolved in the water which are not very favorable to the survival of these germs [19]-[21].

C. Environmental risks of bacteriological parameters

Classical indicator bacteria have proved to be useful in studies describing the environmental distribution of waste water discharges [22]-[29]. Pathogen contamination of aquatic ecosystems is known to occur from a range of sources including municipal waste water effluents, agricultural wastes, and wildlife [30]. The World Health Organization (WHO) has stated that infectious diseases are the world's single largest source of human mortality [31]. Many of these infectious diseases are waterborne and have tremendous adverse impacts in developing countries. Fecal Coliforms are



bacteria found in the digestive tract of warm-blooded animals. The level of coliforms and streptococci in the environment is an indicator of other contaminants. Shellfish can become contaminated through filtering fecal coliforms from the water during feeding, which can make humans sick if the shellfish are eaten. The studied bacteria are excellent indicators to confirm the quality and the level of biological contamination of waters of Bouregreg estuary. The results of the bacteriological analysis showed high levels of fecal coliforms in water samples collected at the station S3 in comparison with the other stations. This is explained by the emplacement of this station which is located next to a main rejection of Oued Akrach and by the conditions and the elements necessary for the survival of these bacteria in the studied waters. Indeed, the waters of this station are of mixed origin with human dominance, In addition to that this station is located in upstream sites which are characterized by stagnant waters where maritime influence is very weak.

#### IV. CONCLUSION

The study area (Bouregreg estuary) is an area of natural (continental and marine) and anthropogenic (liquid waste and / or solid industrial and domestic). From the economic and social plan, the collected data from this research will allow better management of the protection of the coastal marine environment, to the benefit of the development of various riverine activities, notably fishing, tourism, swimming and water sports. The need for the installation of sewage treatment plants in the study area is of major and immediate importance in order to improve the quality of fluvial waters on the one hand and to contribute to the improvement of the state of the coastal marine environment on the other hand.

#### REFERENCES

[1] M. Cheggour, A. Chafik, N.S. Fisher, S. Benbrahim. Metal concentrations in sediments and clams in four Moroccan estuaries. *Mar. Environ. Res.*, 59, (2005) 119-137.

[2] L. Tahiri, L. Bannasser, L. Idrissi, M. Fekhaoui, A. El Abidi, A. Mouradi, Contamination métallique de *Mytilus galloprovincialis* et des sédiments au niveau de l'estuaire de Bouregreg (Maroc), *Water qual. Res. J.*, 40, 1, (2005) 111-119.

[3] L.M. Bannasser, M. Fekhaoui, O. Mameli. Assessment of the metallic contamination of the low Sebou sediments. *Ann. Chimica*, 90, (2000) 637-644.

[4] R. Bchitou, A. Bouhaouss, A. Foutlane. The exchange of phosphorus and nitrogen ions in water sediment interface in SebouRiver (Morocco). *Phys. Chem. News*, 31, (2006) 99-105.

[5] G.K. Elmund, M.J. Allen, E.W. Rice. Comparison of *Escherichia coli*, total coliform and fecal coliform populations as indicators of wastewater treatment efficiency. *Water Environ. Res.*, 71, (1999) 332-339.

[6] E. Emmanuel, K. Thélys, M. Mompoin, J.M. Blanchard, Y. Perrodin. Evaluation des dangers environnementaux liés au rejet des eaux usées urbaines dans la Baie de Port-au-Prince en Haïti Submitted: Livre « Eau et Environnement » du Réseau « Droit de l'Environnement » de l'Agence Universitaire de la francophonie (AUF). Port-au- Prince, (2004) pp.15.

[7] AFNOR (Agence Française de Normalisation). *Eaux-méthodes d'essai*. In : Recueil de normes françaises (6ème édition). La Défense, Paris, 2001.

[8] CEAEQ (Centre d'Expertise en Analyse Environnementale du Québec). Recherche et dénombrement des entérocoques: méthode par filtration sur membrane, MA. 700 – Ent 1.0, Rév. 3, Ministère du Développement durable, de l'Environnement et des Parcs du Québec, pp.23, 2006.

[9] E.M. Clausen, B.L. Green, W. Litsky. Fecal streptococci: indicators of pollution. Dans: Hoadley, AW et BJ Dutka, édit., *Bacterial Indicators/Health hazards associated with water*. Am. Soc. Test. Mat. (ASTM), 635, (1977) 247-264.

[10] J.A.E. Farrow, J. Kruze, B.A. Phillips, A.J. Bramley, M.D. Collins. Taxonomic studies of *S. bovis* and *S. equinus*: description of *S. alactolyticus* sp. no. and *S. saccharolyticus* sp. nov. *System. Appl. Microbiol.*, 5, (1984) 467-482.

[11] G. Bitton. *Wastewater microbiology*. 2e éd. John Wiley & Sons, NY, pp.578, 1999.

[12] C. Gleeson, N. Gray. *The coliform index and waterborne disease*. E & FN Spoon, pp. 194, 1997.

[13] E.E. Geldreich, B.A. Kenner. Concepts of fecal streptococci in stream pollution. *J. Water Poll. Cont. Fed.* 41, (1969) R336-R341.

[14] Masalha, Mahmud et al. Analysis of Transcription of the *Staphylococcus Aureus* Aerobic Class Ib and Anaerobic Class III Ribonucleotide Reductase Genes in Response to Oxygen. *J. Bacteriol.*, 183.24 (2001): 7260–7272.

[15] R. Baylet, G. Mandin. Lagunage et virologie des eaux usées. *La technique de l'eau et de l'assainissement*, 383, (1978) 19-22

[16] P. Boutin. Implication sanitaire de l'assainissement des petites collectivités et l'assainissement autonome. XVIème journée de l'hydraulique, (Nantes) , (1982) 14-15 et 16.

[17] ONEP (Office National de l'Eau potable). *Caractérisation quantitative et qualitative des eaux usées (Maroc)*, 1999.

[18] WHO (World Health Organization). *Use of wastewater in agriculture and aquaculture*. Technical Report Series 778, 1989.

[19] F. Benmessaoud. *Qualité physico-chimique, métallique et bactériologique des eaux de l'estuaire du Bou Regreg et impact sur la biologie et la démographie de *Venerupis decussata* (LINNE, 1758) et *Cardium edule* (LINNE, 1767)*, Université Mohammed V- Agdal, Faculté des Sciences Rabat, 2007.

[20] M. EL AMRAOUI, Y. SALAMA, B. EL AMRAOUI, N. LAZRAK, M. MOUNKAD, Evaluation of the physicochemical and some trace elements of the surface waters of the moroccan atlantic estuary: case of the estuary of the river Bouregreg, *Carpath J Earth Env*, 10 (2), (2015) 189 – 198.

[21] N. Zerki, R. Bchitou, A. Bouhaouss. Characterization of some trace elements in natural waters of the bouregreg estuary (rabat, morocco) by a chemometric study, *Phys. Chem. News*, 59, (2011) 121-126.

[22] C.M. Davies, J.A.H. Long, M. Donald, N.J. Ashbolt. Survival of fecal microorganisms in marine and freshwater sediments. *Appl Environ Microbiol.*, 61, (1995) 1888–1896.

[23] E.L. Elliott, R.R. Colwell. Indicator organisms for estuarine and marine waters. *FEMS Microbiol Rev.*, 32, (1985) 61–79.

[24] D.J. Grimes, F.L. Singleton, J. Stemmler, L. Palmer, P. Brayton, R.R. Colwell. Microbiological effects of wastewater effluent discharged into coastal waters of Puerto Rico. *Water Res.*, 18, (1984) 613–619.

[25] J.P. Howington, G.A. McFeters, J.P. Barry, J.J. Smith. Distribution of the McMurdo Station sewage plume. *Mar Pollut Bull.*, 25, (1992) 324–327.

[26] G.A. McFeters, J.P. Barry, J.P. Howington. Distribution of enteric bacteria in Antarctic seawater surrounding a sewage outfall. *Water Res.*, 27, (1993) 645–650.

[27] J.H. Paul, J.B. Rose, S. Jaing, C. Kellogg, E.A. Shinn. Occurrence of fecal indicator bacteria in surface waters and the subsurface aquifer in Key Largo, Florida. *Appl Environ Microbiol.*, 61, (1995) 2235–2241.

[28] D.L. Sorensen, S.G. Eberl, R.A. Dicksa. *Clostridium perfringens* as a point source in nonpoint polluted streams. *Water Res.*, 23, (1989) 191–197.

[29] M. Takizawa, W.L. Straube, R.T. Hill, R.R. Colwell. Near-bottom pelagic bacteria at deep-water sewage sludge disposal site. *Appl Environ Microbiol.*, 59, (1993) 3406–3410.

[30] Environment Canada. *Threats to Sources of Drinking Water and Aquatic Ecosystem Health in Canada*. National Water Research Institute, Burlington, Ontario. NWRI Scientific Assessment Report Series No.1. pp.72, 2001.

[31] WHO (World Health Organization), , *World health report: fighting disease, fostering development*. World Health Organization, Geneva, 8, (1996) 617–645.